

**IN VITRO PROPAGATION OF CHAMAECYPARIS OBTUSA SIEB. ET ZUCC.**

**Ji Yun Min<sup>1</sup>, Dong Jin Park<sup>2</sup>, Mi Jin Jeong<sup>2</sup>, Hyun Jin Song<sup>2</sup>, Yong Duck Kim<sup>3</sup>, Young Min Kang<sup>4</sup>,  
Chandrakant Shivappa Karigar<sup>5</sup>, and Myung Suk Choi<sup>2\*</sup>**

<sup>1</sup>Sancheong Oriental Medicinal Herb Institute, Sancheong-gun, Gyeongnam, 666-831, South Korea

<sup>2</sup>Division of Environmental Forest Science, Gyeongsang National University, Jinju, 660-701, South Korea, \*Fax: + 82-55-753-6015, \*E-mail: mschoi@gnu.ac.kr

<sup>3</sup>Institute of Hadong Green Tea, Hadong-gun, Gyeongnam, 667-804, South Korea

<sup>4</sup>Department of Forestry, Mississippi State University, P. O. Box 9681, MS39762, USA

<sup>5</sup>Division of Biochemistry, Department of Chemistry, Central College Campus, Bangalore University, Bangalore, 560-001, India

**REFERENCES**

- AMANO T., SAKATANI M. (1987). Vegetative propagation by tissue culture of hinoki (*Chamaecyparis obtusa*). I. Plantlet formation from cotyledon explants. Bulletin of the Nara Forest Experiment Station, 17: 19-25.
- GAMBORG O. L., MILLER R. A., OJIMAKI K. (1968). Nutrient requirements of suspension cultures of soybean root cells. Experimental Cell Research, 50: 51-158.
- GEORGE E. F., SHERRINGTON P. D. (1984). Plant propagation by tissue culture. Handbook and Directory of Commercial laboratories. Eversley, U.K.: Exegetics Ltd. VIII+709 pp.
- GOULD A. R. (1986). Factors controlling generation of variability *in vitro*. In: Vasil I. K. (Ed.). Cell culture and somatic cell genetics in plants. Academic Press, Orlando: 549-567.
- IDE Y., YAMAMOTO S. (1996). Adventitious root formation on *in vitro* micro cuttings of hinoki. Journal of the Japanese Forest Society, 68: 296-298.
- ISHII K. (1986). *In vitro* plantlet formation from adventitious buds on juvenile seedlings of Hinoki cypress (*Chamaecyparis obtusa*). Plant Cell, Tissue and Organ Culture, 7: 247-255.
- ISHII K. (2002). Liquid culture and transformation of hinoki cypress (*Chamaecyparis obtusa* Sieb. et Zucc.). Journal of Forest Research, 7: 99-104.
- ISHII K., MARUYAMA E., HOSOI Y. (2003). Plant regeneration by somatic embryogenesis from *in vitro*-cultured shoots of hinoki cypress (*Chamaecyparis obtusa* Sieb. et Zucc.). Propagation of Ornamental Plants, 3: 19-22.
- ISHII K., SATO T. (1989). *In vitro* plantlet regeneration in some species of Japanese conifers. Proceedings of 6th International Congress, SABRAO: 869-872.
- JEONG M. J., SONG H. J., PARK D. J., MIN J. Y., JO J. S., KIM B. M., KIM H. G., KIM Y. D., KIM R. M., KARIGAR C. S., CHOI M. S. (2009). High frequency plant regeneration following abnormal shoot organogenesis in the medicinal tree *Hovenia dulcis*. Plant Cell Tissue and Organ Culture, 98: 59-65.
- KATASE M. (1993). Factors affecting proliferation of shoot in mulberry axillary bud culture. The Journal of Sericultural Science of Japan, 62: 152-161.
- LINSMAIER E. M., SKOOG F. (1965). Organic growth factor requirements of tobacco tissue cultures. Physiologia Plantarum, 18: 100-127.
- LLOYD G. B., McCOWN B. H. (1981). Commercially feasible micropropagation of mountain laurel (*Kalmia laticolia*) by use of shoot tip culture. Proceedings of the International Plant Propagators Society, 30: 421-437.
- LU M. C. (2002). Micropropagation of *Morus latifolia* Poilet using axillary buds from mature trees. Scientia Horticulturae, 96: 329-341.
- MARUYAMA E., HOSOI Y., ISHII K. (2003). Somatic embryo culture for propagation, artificial seed production, and conservation of sawaracypress (*Chamaecyparis pisifera* Sieb. et Zucc.) Journal Forest Research, 8: 1-8.
- MORTE M. A., HONRUBIA M., PIQUERAS A. (1992). Micropropagation of *Tetraclinis articulata* (Vahl) masters (Cupressaceae). Plant Cell, Tissue and Organ Culture, 28: 231-233.
- MURASHIGE T., SKOOG F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue culture. Physiologia Plantarum, 15: 473-497.
- NITSCH J. P., NITSCH C. (1969). Haploid plants from pollen grains. Science, 163: 85-87.
- ONO M., ASAI T., WATANABE H. (1998). Hinokitiol production in a suspension culture of *Calocedrus formosana* Florin. Bioscience Biotechnology and Biochemistry, 62: 1653-1659.
- QUIRIN M., LEPOIVRE P. (1977). Improved media for *in vitro* culture of *Prunus* sp. Acta Horticulturae, 78: 437-442.
- RATHORE J. S., RATHORE V., SHEKHAWAT N. S., SINGH R. P., LILER G., PHULWARIA M., DAGLA H. R. (2004). Micropropagation of woody plants. In: Srivastava P. S., Narula A., Srivastava S. (Eds). Plant Biotechnology and Molecular Markers: 195-205.
- SAS INSTITUTE INC (2001). Using the SAS® System Release 6.12 on Windows™, Gary Mehler, SAS Institute Inc., Cary, NC: 1-9 pp.
- SCHENK R. U., HILDEBRANDT A. C. (1972). Medium and techniques for induction and growth of monocotyledonous and dicotyledonous plant cell cultures. Canadian Journal of Botany, 50: 199-204.
- TANIGUCHI T., KURITA M., ITAHANA N., KONDO T. (2004). Somatic embryogenesis and plant regeneration from immature zygotic embryos of Hinoki cypress (*Chamaecyparis obtusa* Sieb. et Zucc.). Plant Cell Reports, 23: 26-31.
- THORPE T. A., HARRY I. S., KUMAR P. P. (1991). Application of micropropagation to forestry. In: Debergh P. C., Zimmerman R. H. (Eds). Micropropagation: Technology and Application. Kluwer Academic, Dordrecht: 311-316.
- WHITE P. R. (1963). The cultivation of animal and plant cells. 2nd edition. The Ronald Press Co, New York, 228 pp.