

IN VITRO GROWTH OF *ALOE BARBADENSIS* MILL.: THE EFFECT OF ACTIVATED CHARCOAL ON MEDIUM pH, NITROGEN UPTAKE AND ELEMENTS CONTENT OF SHOOTS

**Daniela Borgognone¹, Andrea Marcucci², Giuseppe Colla¹, Marco Renzaglia²,
Elvira Rea², and Mariateresa Cardarelli^{2*}**

¹Department of Agriculture, Forestry, Nature and Energy, University of Tuscia,
via San Camillo De Lellis, 01100 Viterbo, Italy

²Agricultural Research Council - Research Centre for the Soil-Plant System, 00184 Roma, Italy,

*Fax: + 39-(0) 6-7005711, *E-mail: mteresa.cardarelli@entecra.it

REFERENCES

- ADELBERG J. W., DELGADO M. P., TOMKINS J. T. (2010). Spent medium analysis for liquid culture micropropagation of *Hemerocallis* on Murashige and Skoog medium. *In Vitro Cellular & Developmental Biology-Plant*, 46: 95-107.
- AHLAWAT K. S., KHATKAR B. S. (2011). Processing, food applications, and safety of *Aloe vera* products: a review. *Journal of Food Science and Technology*, 48: 525-533.
- BORGOGNONE D., COLLA G., CARDARELLI M. (2010). Coltura *in vitro* di *Aloe barbadensis*: ruolo del carbone attivo nelle fasi di allungamento ed ambientamento di germogli micro propagati. *Acta Italus Hortus*, 1: 43-47.
- BREMNER J. M. (1965). Total nitrogen. *In*: Black C. A., Evans D. D., White I. L., Ensminger L. E., Clark F. E. (Eds). *Methods of soil analysis*. Agronomy Monograph 9, Part 2: 1149-1178.
- BRITO G., COSTA E. A., COELHO E. C., SANTOS C. (2009). Large-scale field acclimatization of *Olea maderensis* micropropagated plants: morphological and physiological survey. *Trees*, 23: 1019-1031.
- CATALDO D. A., HAROON M., SCHRADER L. E., YOUNGS V. L. (1975). Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Communications in Soil Science and Plant Analysis*, 6: 71-80.
- CHEN F. Q., FU Y., WANG D. I., GAO X., WANG I. (2007). The effect of plant growth regulators and sucrose on the micropropagation and microtuberization of *Dioscorea nipponica* Makino. *Journal of Plant Growth Regulation*, 26: 38-45.
- DE OLIVEIRA E. T., CROCOMO O. J., FARINHA T. B., GALLO L. A. (2009). Large-scale micropropagation of *Aloe vera*. *HortScience*, 44: 1675-1678.
- DUMAS E., MONTEUUIS O. (1995). *In vitro* rooting of micropropagated shoots from juvenile and mature *Pinus pinaster* explants: influence of activated charcoal. *Plant Cell, Tissue and Organ Culture*, 40: 231-235.
- EYMAR E., ALEGRE J., TORIBIO M., LÓPEZ-VELA D. (2000). Effect of activated charcoal and 6-benzyladenine on *in vitro* nitrogen uptake by *Lagerstroemia indica*. *Plant Cell, Tissue and Organ Culture*, 63: 57-65.
- FIELD A. (2005). *Discovering statistics using SPSS for windows*, 2nd edn. SAGE Publications Ltd., London, 816 pp.
- FRIDBORG G., PEDERSÉN M., LANDSTRÖM L., ERIKSSON T. (1978). The effect of activated charcoal on tissue cultures: adsorption of metabolites inhibiting morphogenesis. *Physiologia Plantarum*, 43: 104-106.
- GUBBUK H., PEKMEZCI M. (2006). *In vitro* propagation of banana (*Musa* spp.) using thidiazuron and activated charcoal. *Acta Agriculturae Scandinavica*, 56: 65-69.
- HASHEM ABADI D., KAVIANI B. (2010). *In vitro* proliferation of an important medicinal plant *Aloe* - a method for rapid production. *Australian Journal of Crop Science*, 4: 216-222.
- HAZARIKA B. N. (2006). Morpho-physiological disorders in *in vitro* culture of plants. *Scientia Horticulturae*, 108: 105-120.
- HEMPHILL J. K., MAIER C. G. A., CHAPMAN K. D. (1998). Rapid *in-vitro* plant regeneration of cotton (*Gossypium hirsutum* L.) *Plant Cell Report*, 17: 273-278.
- KARLA Y. P. (1998). *Handbook of reference methods for plant analysis*. CRC Press Inc, Boca Raton, FL: 165-170.
- KEIJZER C. J., CRESTI M. (1987). A comparison of anther tissue development in male sterile *Aloe vera* and male fertile *Aloe ciliaris*. *Annals of Botany*, 59: 533-542.
- KET N. V., HAHN E. J., PARK S. Y., CHAKRABARTY D., PAEK K. Y. (2004). Micropropagation of an endangered orchid *Anoectochilus formosanus*. *Biologia Plantarum*, 48: 339-344.
- KRISHNA H., SAIRAM R. K., SINGH S. K., PATEL V. B., SHARMA R. R., GROVER M., NAIN L., SACHDEV A. (2008). Mango explant browning: effect of ontogenic age, mycorrhization and pre-treatments. *Scientia Horticulturae*, 118: 132-138.
- MALDA G., BACKHAUS R. A., MARTIN C. (1999). Alterations in growth and crassulacean acid metabolism (CAM) activity of *in vitro* cultured cactus. *Plant Cell, Tissue and Organ Culture*, 58: 1-9.
- MURASHIGE T., SKOOG F. (1962). A revised medium for rapid growth and bioassay with tobacco tissues cultures. *Physiologia Plantarum*, 15: 473-497.
- NATALI L., SANCHEZ I. C., CAVALLINI A. (1990). *In vitro* culture of *Aloe barbadensis* Mill.: Micropropagation from vegetative meristems. *Plant Cell, Tissue and Organ Culture*, 20: 71-74.
- NURUDDIN A. A., CHANG M. (1999). Responses of herbaceous mimosa (*Mimosa strigillosa*), a new reclamation species to soil pH. *Resources Conservation and Recycling*, 27: 287-298.

- OWEN H. R., WENGERD D., MILLER A. R. (1991). Culture medium pH is influenced by basal medium, carbohydrate source, gelling agent, activated charcoal, and medium storage method. *Plant Cell Reports*, 10: 583-586.
- PAEK K. Y., YEUNG E. C. (1991). The effects of 1-naphthaleneacetic acid and N⁶-benzyladenine on the growth of *Cymbidium forrestii* rhizomes *in vitro*. *Plant Cell, Tissue and Organ Culture*, 24: 65-71.
- PAN M. J., VAN STADEN J. (1998). The use of charcoal in *in vitro* culture - a review. *Plant Growth Regulation*, 26: 155-163.
- PAN M. J., VAN STADEN J. (1999). Effect of activated charcoal, autoclaving and culture media on sucrose hydrolysis. *Plant Growth Regulation*, 29: 135-141.
- PEUKE A. D., TISCHNER R. (1991). Nitrate uptake and reduction of aseptically cultivated spruce seedlings, *Picea abies* (L.) Karst. *Journal of Experimental Botany*, 42: 723-728.
- QUOIRIN M., DA SILVA M. C., MARTINS K. G., DE OLIVEIRA D. E. (2001). Multiplication of juvenile black wattle by microcuttings. *Plant Cell, Tissue and Organ Culture*, 66: 199-205.
- RAMIREZ-MALAGON R., BORODANENKO A., BARRERA-GUERRA J. L., OCHOA-ALEJO N. (2001). Shoot number and shoot size as affected by growth regulators in *in vitro* cultures of *Spathiphyllum floribundum* L. *Scientia Horticulturae*, 89: 227-236.
- RUAN J., GERENDÁS J., HÄRDTER R., SATTLEMACHER B. (2007). Effect of nitrogen form and root-zone pH on growth and nitrogen uptake of Tea (*Camellia sinensis*) plants. *Annals of Botany*, 99: 301-310.
- RUSSOWSKI D., MAURMANN N., RECH S. B., FETT-NETO A. G. (2006). Role of light and medium composition on growth and valepotriate contents in *Valeriana glechomifolia* whole plant liquid cultures. *Plant Cell, Tissue and Organ Culture*, 86: 211-218.
- SHIBLI R. A., MOHAMMAD M. J., AJLOUNI M. M., SHATNAWI M. A., OBEIDAT A. F. (1999). Stability of chemical parameters of tissue culture medium (pH, osmolarity, electrical conductivity) as a function of time of growth. *Journal of Plant Nutrition*, 22: 501-510.
- THIND S. K., JAIN N., GOSAL S. S. (2008). Micropropagation of *Aloe vera* L. and estimation of potentially active secondary constituents. *Phytomorphology: An International Journal of Plant Morphology*, 58: 65-71.
- THOMAS T. D. (2008). The role of activated charcoal in plant tissue culture. *Biotechnology Advances*, 26: 618-631.
- VAN WINKLE S. C., JOHNSON S., PULLMAN G. S. (2003). The impact of gelrite and activated carbon on the elemental composition of two conifer embryogenic tissue initiation media. *Plant Cell Reports*, 21: 1175-1182.
- VAN WINKLE S. C., PULLMAN G. S. (2003). The combined impact of pH and activated carbon on the elemental composition of a liquid conifer embryogenic tissue initiation medium. *Plant Cell Reports*, 22: 303-311.
- VAN WINKLE S. C., PULLMAN G. S. (2005). Achieving desired plant growth regulator levels in liquid plant tissue culture media that include activated carbon. *Plant Cell Reports*, 24: 201-208.
- WANN S. R., VEAZEY R. L., KAPHAMMER J. (1997). Activated charcoal does not catalyze sucrose hydrolysis in tissue culture media during autoclaving. *Plant Cell, Tissue and Organ Culture*, 50: 221-224.
- WEATHERBURN M. W. (1967). Phenol hypochlorite reaction for determination of ammonia. *Analytical Chemistry*, 39: 917-920
- WU P., HU B., LIAO C. Y., ZHU J. M., WU Y. R., SENADHIRA D., PATERSON A. H. (1998). Characterization of tissue tolerance to iron by molecular markers in different lines of rice. *Plant and Soil*, 203: 217-226.
- YAN N., HU H., HUANG J., XU K., WANG H., ZHOU Z. (2006). Micropropagation of *Cypripedium flavum* through multiple shoots of seedlings derived from mature seeds. *Plant Cell, Tissue and Organ Culture*, 84: 113-117.