

**EFFECTS OF NITROGEN SOURCE AND WAVELENGTH OF LED-LIGHT ON
ORGANOGENESIS FROM LEAF AND SHOOT TIP CULTURES
IN *LYSIONOTUS PAUCIFLORUS* MAXIM.**

Yuan-Xue Lu¹, Toshinari Godo², Kazuhiro Fujiwara³, Kai-Yuan Guan¹, and Masahiro Mii^{4*}

¹Key Laboratory of Economic Plants and Biotechnology, Kunming Institute of Botany, Chinese Academy of Sciences, 132 Lanhei Road, Kunming, Yunnan 650204, China

²Botanic Gardens of Toyama, 42 Kamikutsuwada, Fuchu-machi, Toyama 939-2713, Japan

³Department of Biological and Environmental Engineering, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Bunkyo-ku, Tokyo, 113-8657, Japan

⁴Laboratory of Plant Cell Technology, Graduated School of Horticulture, Chiba University, 648 Matsudo, Matsudo, Chiba 271-8510, Japan, * Fax: +81 4-7134-8008,

*E-mail: miim@faculty.chiba-u.jp

REFERENCES

- BAQUE M. A., HAHN E. J., PAEK K. Y. (2010). Induction mechanism of adventitious root from leaf explants of *Morinda citrifolia* as affected by auxin and light quality. *In Vitro Cellular & Developmental Biology-Plant*, 46: 71-80.
- BAQUE M. A., SHIN Y. K., ELSHMARI T., LEE E. J., PAEK K. Y. (2011). Effect of light quality, sucrose and coconut water concentration on the micropropagation of *Calanthe* hybrids ('Bukduseong' × 'Hyesung' and 'Chunkwang' × 'Hyesung'). *Australian Journal of Crop Science*, 5: 1247-1254.
- BHOJWANI S. S., RAZDAN M. K. (1996). Tissue culture media. *In: Bhojwani S. S., Razdan M. K. (Eds). Plant Tissue Culture: Theory and Practice, A Revised Edition, Elsevier, Amsterdam: 39-62.*
- CYBULARZ-URBAN T., HANUS-FAJERSKA E., ŚWIDERSKI A. (2007). Effect of light wavelength on *in vitro* organogenesis of a *Cattleya* hybrid. *Acta Biologica Cracoviensia, Series Botanica*, 49/1: 113-118.
- FASOLO F., ZIMMERMAN R. H., FORDHAM I. (1989). Adventitious shoot formation on excised leaves of *in vitro* grown shoots of apple cultivars. *Plant Cell, Tissue and Organ Culture*, 16: 75-87.
- GODO T., LU Y. X., MII M. (2010). Micropropagation of *Lysionotus pauciflorus* Maxim. *In: Jain S. M., Ochatt S. J. (Eds). Protocols for In Vitro Propagation of Ornamental Plants, Methods in Molecular Biology, Helsinki, Finland, Humana Press, 589: 127-139.*
- GULATI A., JAIWAL P. K. (1994). Plant regeneration from cotyledonary node explants of mungbean (*Vigna radiata* (L.) Wilczek). *Plant Cell Reports*, 13: 523-527.
- HALPERIN W., WETHRELL D. (1965). Ammonium requirement for embryogenesis *in vitro*. *Nature*, 205: 519-520.
- HARUKI K., INAMURA H. (2003). Propagation and growing of *Lysionotus pauciflorus* Maxim. *Bulletin of the Shimane Agricultural Experimental Station*, 34: 61-70.
- HASSAN A. Q., SKIRVIN R. M., BELOW F. E. (1991). Influence of nitrogen form and NH₄⁺-N:NO₃⁻-N ratios on adventitious shoot formation from pear (*Pyrus communis*) leaf explants *in vitro*. *Plant Cell, Tissue and Organ Culture*, 27: 315-319.
- HUNTER D. C., BURRITT D. J. (2004). Light quality influences adventitious shoot production from cotyledon explants of lettuce (*Lactuca sativa* L.). *In Vitro Cellular & Developmental Biology-Plant*, 40: 215-220.
- IVANOVA M., VAN STADEN J. (2009). Nitrogen source, concentration, and NH₄⁺: NO₃⁻ ratio influence shoot regeneration and hyperhydricity in tissue cultured *Aloe polyphylla*. *Plant Cell, Tissue and Organ Culture*, 99: 167-174.
- KIM S. J., HAHN E. J., HEO J. W., PAEK K. Y. (2004). Effects of LEDs on net photosynthetic rate, growth and leaf stomata of *Chrysanthemum* plantlets *in vitro*. *Scientia Horticulturae*, 101: 143-151.
- LIAN M. L., MURTHY H. N., PAEK K. Y. (2002). Effects of light emitting diodes (LEDs) on the *in vitro* induction and growth of bulblets of *Lilium* Oriental hybrid 'Pesaro'. *Scientia Horticulturae*, 94: 365-370.
- LIU Y., WAGNER H., BAUER R. (1996). Nevadensin glycosides from *Lysionotus pauciflorus*. *Phytochemistry*, 42: 1203-1205.
- LIU Y., WAGNER H., BAUER R. (1998). Phenylpropanoids and flavonoid glycosides from *Lysionotus pauciflorus*. *Phytochemistry*, 46: 339-343.
- LU Y. X., GODO T., CHIN D. P., MII M., GUAN K. Y. (2006). Establishment of callus culture with high regeneration ability from leaf segments of *Lysionotus pauciflorus* Maxim. *Propagation of Ornamental Plants*, 6: 180-186.
- MOREIRA DA SILVA M. H., DEBERGH P. C. (1997). The effect of light quality on the morphogenesis of *in vitro* cultures of *Azorina vadalii* (Wats.) Feer. *Plant Cell, Tissue and Organ Culture*, 51: 187-193.
- MURASHIGE T., SKOOG F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- NAGAKUBO T., NAGASAWA A., OHKAWA H. (1993). Micropropagation of garlic through *in vitro* bulblet formation. *Plant Cell, Tissue and Organ Culture*, 32: 175-183.
- POUDEL P. R., KATAOKA I., MOCHIOKA R. (2008). Effect of red- and blue-light-emitting diodes on growth and morphogenesis of grapes. *Plant Cell, Tissue and Organ Culture*, 92: 147-153.

- PREDIERI S., FASOLO F., MALAVASI F. (1989). High-frequency shoot regeneration from leaves of the apple rootstock M26 (*Malus pumila* Mill.). *Plant Cell, Tissue and Organ Culture*, 17: 133-142.
- SHIN K. S., MURTHY H. N., HEO J. W., HAHN E. J., PAEK K. Y. (2008). The effect of light quality on the growth and development of *in vitro* cultured *Doritaenopsis* plants. *Acta Physiologiae Plantarum*, 30: 339-343.
- TSAI C. J., SAUNDERS J. W. (1999). Evaluation of sole nitrogen sources for shoot and leaf disc cultures of sugarbeet. *Plant Cell, Tissue and Organ Culture*, 59: 47-56.