

SHORTENING OF SELECTION TIME OF *ROSA HYBRIDA* BY *IN VITRO* CULTURE OF ISOLATED EMBRYOS AND IMMATURE SEEDS

Matteo Caser¹, Fulvio Dente², Gian Guido Ghione³, Andrea Mansuino³,
Annalisa Giovannini², and Valentina Scariot^{1*}

¹Department of Agricultural, Forest and Food Sciences, University of Torino, 2 Paolo Braccini square,
10095 Grugliasco (TO), Italy.

²CRA-FSO, Ornamental Species Research Unit, 508 Corso degli Inglesi str., 18038 Sanremo (IM), Italy,
*Fax +39 011 6708798, *E-mail valentina.scariot@unito.it

³NIRP International, Az. Agricola di Ghione L. & Figli, 1 San Rocco str., Fraz. Bevera,
18039 Ventimiglia (IM), Italy.

REFERENCES

- ARENE L., PELLEGRINO C., GUDIN S. (1993). A comparison of the somaclonal variation level of *Rosa hybrida* L. cv. Meirutral plants regenerated from callus or direct induction from different vegetative and embryonic tissues. *Euphytica*, 71: 83-90.
- ARUNACHALAM V., KAICKER U. S. (1994). *In vitro* germination. A potential commercial method for roses. In: PRAKASH J., BHANDRY K. R. (Eds). *Floriculture, technology, trades and trends*. Oxford and IBH publishing CO. PVT. LTD, New Delhi: 410-412.
- ASEN S. (1948). Embryo culture of rose seeds. *American Rose Annual*, 33: 151-153.
- ASEN S., LARSEN R. E. (1951). Artificial culturing of rose embryos. Pennsylvania State University. Agricultural Experiment Station Progress Report, 40: 1-4.
- BILLINGS W. D., MOONEY H. A. (1968). The ecology of arctic and alpine plants. *Biological Reviews*, 43: 481-529.
- BOSCO R., CASER M., GHIONE G. G., MANSUINO A., GIOVANNINI A., SCARIOT V. (2014). Dynamics of abscisic acid and indole-3-acetic acid during the early-middle stage of seed development in *Rosa hybrida*. *Plant Growth Regulation*, DOI 10.1007/s10725-014-9950-8
- BOUSEWINKEL F. D., BOUMAN F. (1995). The seed: Structure and function. In: KIGEL J., GALILI G. (Eds). *Seed Development and Germination*, Marcel Dekker, Inc. New York: 1-24.
- BRESSAN P. H., KIM Y. J., HYNDMAN S. E., HASEGAWA P. M., BRESSAN R. A. (1982). Factors affecting *in vitro* propagation of rose. *Journal of American Society for Horticultural Science*, 107: 979-990.
- BURGER D. W., LIU L., ZARY K. W., LEE C. I. (1990). Organogenesis and plant regeneration from immature embryos of *Rosa hybrida* L. *Plant Cell, Tissue and Organ Culture*, 21: 147-152.
- CARELLI B. P., ECHEVERRIGARAY S. (2002). An improved system for the *in vitro* propagation of rose cultivars. *Scientia Horticulturae*, 92: 69-74.
- CZABATOR F. J. (1962). Germination value: an index combining speed and completeness of pine seed germination. *Forest Science*, 8: 386-396.
- DE COCK K., SCARIOT V., LEUS L., DE RIEK J., VAN HUYLENBROECK J. (2007). Understanding genetic relationships of wild and cultivated rose and the use of species in breeding. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources*, 2 (52), 10 pp.
- DE VRIES D. P., DUBOIS L. A. M. (1987). The effect of temperature on fruit-set, seed set and seed-germination in Sonia × Hadley hybrid tea-rose crosses. *Euphytica*, 36: 117-120.
- GUDIN S. (1994). Embryo rescue in *Rosa hybrida* L. *Euphytica*, 72: 205-212.
- GUDIN S. (2000). Rose: genetics and breeding. *Plant Breeding Reviews*, 17: 59-189.
- GUDIN S., ARENE L., CHAVAGNAT A., BULARD C. (1990). Influence of endocarp thickness on rose achene germination: genetic and environmental factors. *HortScience*, 25: 786-788.
- GUZICKA M., ZIELIŃSKI J., TOMASZEWSKI D., GAWLAK M. (2012). Anatomical study on the developing pericarp of selected *Rosa* species (Rosaceae). *Dendrobiology*, 68: 77-87.
- HÜBNER S. (2012). International statistics. *Flowers and Plants 2012*. AIPH/Union Fleurs, 60, 142 pp.
- ISTA. (2014). International Rules for Seed Testing 2014. Ordinary General Meeting 2013, Antalya, Turkey. Online ISSN 2310-3655 <http://www.seedtest.org/seedhealthmethods>
- KELLY K. M., VAN STADEN J., BELL W. E. (1992). Seed coat structure and dormancy. *Plant Growth Regulation*, 11: 201-209.
- KUNITAKE H., IMAMIZO H., MII M. (1993). Somatic embryogenesis and plant regeneration from immature seed-derived calli of rugosa rose (*Rosa rugosa* Thunb.). *Plant Science*, 90: 187-194.
- LAMMERTS W. E. (1946). Use of embryo culture in rose breeding. *Plants and Gardens*, 21 (2): 77-78.
- MARCHANT R., POWER J. B., DAVEY M. R., CHARTIER-HOLLIS J. (1994). Embryo rescue, for the production of F1 hybrids, in English rose. *Euphytica*, 74: 187-193.
- MEYNET J., BARRADE R., DUCLOS A., SIADOUS R. (1994). Dihaploid plants of roses (*Rosa × hybrida* cv. Sonia) obtained by parthenogenesis induced using irradiated pollen and *in vitro* culture of immature seeds. *Agronomie*, 2: 169-175.

- MOHAPATRA A., ROUT G. R. (2005). Study of embryo rescue in floribunda rose. *Plant Cell, Tissue and Organ Culture*, 81: 113-117.
- MURASHIGE T., SKOOG F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- NIKBAKHT A., KAFI M., MIRMASOUMI M., BABALAR M. (2005). Micropropagation of Damask Rose (*Rosa damascena* Mill.) cvs Azaran and Ghamsar. *International Journal of Agriculture & Biology*, 7: 535-538.
- Ogilvie I., Cloutier D., Arnold N., Jui P. Y. (1991). The effect of gibberellic acid on fruit and seed set in crosses of garden and winter hardy *Rosa* accessions. *Euphytica*, 52: 119-123.
- PATI P. K., RATH S. P., SHARMA M., SOOD A., AHUJA P. S. (2006). *In vitro* propagation of rose - a review. *Biotechnology Advances*, 24: 94-114.
- PIPINO L. (2011). Improving seed production efficiency for hybrid rose breeding. PhD thesis, Ghent University and University of Turin, 189 pp.
- PIPINO L., SCARIOT V., GAGGERO L., MANSUINO A., VAN LABEKE M. C., GIOVANNINI A. (2011). Enhancing seed germination in hybrid tea roses. *Propagation of Ornamental Plants*, 11: 111-118.
- PIPINO L., LEUS L., SCARIOT V., VAN LABEKE M. C. (2013). Embryo and hip development in hybrid rose. *Plant Growth Regulation*, 69: 107-116.
- RAM M., PRASAD K. V., KAUR C., SINGH S. K., ARORA A., KUMAR S. (2011). Induction of anthocyanin pigments in callus cultures of *Rosa hybrida* L. in response to sucrose and ammonical nitrogen levels. *Plant Cell, Tissue and Organ Culture*, 104: 171-179.
- ROUT G. R., SAMANTARAY S., MOTTLEY J., DAS P. (1999) Biotechnology of the rose: a review of recent progress. *Scientia Horticulturae*, 81: 201-228.
- SCHEIBER S. M., ROBACKER C. D. (2003). Effect of pericarp removal, gibberellic acid treatment, and stratification on seed germination of *Abelia x grandiflora*. *Journal of Environmental Horticulture*, 21: 34-37.
- SEMENIUK P., STEWART R. N., UHRING J. (1963). Induced secondary dormancy of rose embryos. *American Society of Horticultural Science*, 83: 825-828.
- TILLBERG E. (1983). Levels of endogenous abscisic acid in achenes of *Rosa rugosa* during dormancy release and germination. *Physiologia Plantarum*, 58: 243-248.
- ZHOU Z., BAO W., WU N. (2009). Effects of scarification, stratification and chemical treatments on the germination of *Rosa soulieana* Crépin achenes. In: TAXEIRA DA SILVA J. (Ed.). *Floriculture and Ornamental Biotechnology, Special Issue 1*. Global Science Books, London, UK: 75-80.
- ZLESAK D. C. (2006). Rose: *Rosa × hybrida*. In: ANDERSON N. O. (Ed.). *Flower Breeding and Genetics: Issues, Challenges and Opportunities for the 21st Century*. Springer, Dordrecht, The Netherlands: 695-738.