

IN VITRO PLANT REGENERATION OF ORNAMENTAL CABBAGE
(*BRASSICA OLERACEA* VAR. *LONGATA*)

Yipeng Zhao^{1*}, Jon Green², Miao Yu² and Brian W.W. Grout²

¹Department of Horticulture, Henan Institute of Science and Technology,
Xinxiang, Henan 453003, China, *Fax: + 86-373-304066 *E-mail: yzhao36@hotmail.com
²Postgraduate School, Writtle College, Lordship Road Chelmsford CM1 3RR, United Kingdom

REFERENCES

- Bhalla P. L., Smith N. A. (1998). Comparison of shoot regeneration potential from seedling explants of Australian cauliflower (*Brassica oleracea* var. *botrytis*) varieties. Australian Journal of Agricultural Research, 49 (8): 1261-1266.
- Grout B. W. W. (1990). Meristem tip culture. In: Pollard J. W., Walker J. M. (Eds.). Plant Cell and Tissue Culture. The Humana Press Inc, Clifton, New Jersey: 81-91.
- Grout B. W. W., Crisp P. (1977). Practical aspects of the propagation of cauliflower by meristem culture. Acta Horticulturae, 78: 289-296.
- Grout B. W. W., Crisp P. (1980). The origin and nature of shoots propagated from cauliflower roots. Journal of Horticultural Science, 55 (1): 65-70.
- Grout B. W. W., Westcott R. J., Henshaw G. G. (1977). Scanning electron microscope studies of multiple shoot production by meristem tip cultures of *Solanum x curtilobum*. Annals of Botany, 41: 1113-1116.
- Hwang S. J. (2004). An efficient *in vitro* protocol for propagation of *Zanthoxylum piperitum* DC. Propagation of Ornamental Plants, 2: 37-40.
- Jain S. M. (2001). Tissue culture-derived variation in crop improvement. Euphytica, 118: 153-166.
- Loiseau J., Michaux-Ferrere N., Deunff Y. L. (1998). Histological investigations of somatic embryogenesis in peas. Plant Physiology and Biochemistry, 36: 683-687.
- Murashige T., Skoog F. (1962). A revised medium for rapid rowth and bioassays with tobacco tissue cultures. Physiologia Plantarum, 15: 473-497.
- Msikita W., Skirvin R. M. (1989). *In vitro* regeneration from hypocotyls and seedling cotyledons of tronchuda (*Brassica oleracea* var. *tronchuda* Bailey). Plant Cell, Tissue and Organ Culture, 19: 159-165.
- Narasimhulu S. B., Chopra V. L. (1988). Species-specific shoot regeneration response of cotyledonary explants of Brassicas. Plant Cell Reports, 7: 473-497.
- Pua E. C., Teo S. H., Loh C. S. (1996). Interactive role of ethylene and polyamines on shoot regenerability of Chinese cabbage (*Brassica alboglabra* Bailey) *in vitro*. Journal of Plant Physiology, 149: 138-148.
- Pua E. C., Trinh T. H., Chua N. H. (1989). High frequency plant regeneration from stem explants of *Brassica alboglabra* Bailey *in vitro*. Plant Cell, Tissue and Organ Culture, 17: 143-152.
- Sretenović-Rajčić T., Mijatović M., Vinterhalter D. (2002). *In vitro* culture as a tool for improvement of cabbage cultivars in Yougoslavia. Acta Horticulturae, 579: 209-213.