

CRYOPRESERVATION FOR GENE CONSERVATION OF *ACER MONO MAX*

Young-Goo Park^{1*}, Gi-Sok Kwon² and David Tay³

¹Dept. of Forestry, Kyunpook National University, Deagu 702-701, Republic of Korea,

*Fax: 82-53-950-6708, *E-mail: ygpark@mail.knu.ac.kr

²R&D Institute Tongyang Moolsan Co., Ltd., Yangji-Myeon, Yongin-Si, Gyeonggi-Do 449-825,
Republic of Korea

³Ornamental Plant Germplasm Center, The Ohio State University, 670 Vernon Tharp Street, Columbus, OH
43210-1086, USA

REFERENCES

- Bercetche J., Galerne M., Dereuddre J. (1990). Efficient regeneration of plantlets from embryogenic callus of *Picea abies* (L.) Karst after freezing in liquid nitrogen. *Comptes Rendus de l'Académie des Sciences, Paris* 310: 357-363.
- Dereuddre J., Scottez C., Arnaud Y., Duron M. (1990). Resistance of alginate-coated axillary shoot tips of pear tree (*Pyrus communis* L. cv. *Beurre Hardy*) *in vitro* plantlets to dehydration and subsequent freezing in liquid nitrogen. *Comptes Rendus de l'Académie des Sciences, Paris*, 310: 317-323.
- Ford C. S., Jones N. B., van Staden J. (2000). Cryopreservation and plant regeneration from somatic embryos of *Pinus patula*. *Plant Cell Reports*, 19: 610-615.
- Fukai S., Oe M. (1990). Morphological observation of *Chrysanthemum* shoot tips cultured after cryoprotection and freezing. *Journal of the Japanese Society for Horticultural Science*, 59: 383-387.
- Hirai D., Sakai A. (1999). Cryopreservation of *in vitro*-grown axillary shoot-tip meristems of mint (*Mentha spicata* L.) by encapsulation vitrification. *Plant Cell Reports*, 19: 150-155.
- Kobayashi S., Sakai A., Oiyama I. (1990). Cryopreservation in liquid nitrogen of cultured navel orange (*Citrus sinensis* Osb.) nucellar cells and plant regeneration. *Plant Cell Tissue and Organ Culture*, 23: 15-20.
- Kuoksa T., Hohtola A. (1991). Freeze-preservation of buds from scots pine trees. *Plant Cell Tissue and Organ Culture*, 27: 89-93.
- Langis R., Schnabel B., Earle E. D., Steponkus P. C. (1989). Cryopreservation of *Brassica campestris* L. cell suspensions by vitrification. *CryoLetters*, 10: 421-428.
- Matsumoto T., Sakai A., Nako Y. (1998). A novel preculturing for enhancing the survival of *in vitro*-grown meristems of wasabi (*Wasabia japonica*) cooled to -196°C by vitrification. *CryoLetters*, 19: 27-36.
- Matsumoto T., Sakai A., Yamada K. (1994). Cryopreservation of *in vitro*-grown apical meristems of wasabi (*Wasabia japonica*) by vitrification and subsequent high plant regeneration. *Plant Cell Reports*, 13: 442-226.
- Murashige T., Skoog F. (1962). Revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiologia Plantarum*, 15: 473-497.
- Niino T., Sakai A., Yakuwa H., Nojiri K. (1992). Cryopreservation of *in vitro*-grown shoot tips of apple and pear by vitrification. *Plant Cell Tissue and Organ Culture*, 28: 261-266.
- Niino T., Tashiro K., Suzuki M., Ohuchi S., Magoshi J., Akihama T. (1997). Cryopreservation of *in vitro* grown shoot tips of cherry and sweet cherry by one-step vitrification. *Scientia Horticulture*, 70: 155-163.
- Nishizawa S., Sakai A., Amano Y., Matsuzawa T. (1993). Cryopreservation of asparagus (*Asparagus officinalis* L.) embryogenic suspension cells and subsequent plant regeneration by vitrification. *Plant Science*, 91: 67-73.
- Park Y. G., Son S. H. (1996). Forest genetic conservation based on *in vitro* culture systems. In: Park Y.-G., Sakamoto S. (Eds.). *Biodiversity and Conservation of Plant Genetic Resources in Asia*. Japan Scientific Societies Press, Tokyo: 153-173.
- Paul H., Daigny G., Sangwan, Norreel B. S. (2000). Cryopreservation of apple (*Malus x domestica* Borlch.) shoot tips following encapsulation-dehydration or encapsulation-vitrification. *Plant Cell Reports*, 19: 768-744.
- Phunchindawan M., Hirata K., Sakai A., Miyamoto K. (1997). Cryopreservation of encapsulated shoot primordia induced in horseradish (*Armoracia rusticana*) hairy root cultures. *Plant Cell Reports*, 16: 469-473.

- Reed B. M. (1990). Survival of *in vitro* grown apical meristems of *Pyrus* following cryopreservation. Hort Science, 25: 111-113.
- Reed B. M. (1991). Application of gas-permeable bags for *in vitro* cold storage of strawberry germplasm. Plant Cell Reports, 10: 431-434.
- Sakai A. (1993). Cryogenic strategies for survival of plant cultured cells and meristems cooled to -196°C. Japan International Cooperation Agency, Genetic Resources Project, Reference No.6: 5-30.
- Sakai A. (1995). Cryopreservation of germplasm of woody plants. in Biotechnology In: Bajaj Y. P. S. (Ed.). Agriculture and Forestry. Springer-Verlag Berlin Heidelberg, 32: 53-69.
- Sakai A. (1997). Potentially valuable cryogenic procedure for cryopreservation of cultured plant meristems. In: Razda M. K., Cocking E. C. (Eds.). Conservation of Plant Genetic Resources *in Vitro*. Science Publishers, Inc, Enfield, NH, USA, Vol. 1: 53-66.
- Sakai A., Kobayashi S., Oiyama I. (1990). Cryopreservation of nucellar cells of navel orange (*Citrus sinensis* Osb. var. *brasiliensis* Tanaka) by vitrification. Cryobiology, 27: 657.
- Sakai A., Kobayashi S., Oiyama I. (1991a). Cryopreservation of nucellar cells of navel orange (*Citrus sinensis* Osb.) by a simple freezing method. Plant Science, 54: 243-248.
- Sakai A., Kobayashi S., Oiyama I. (1991b). Survival by vitrification of nucellar cells of navel orange (*Citrus sinensis* var. *brasiliensis* Tanaka) cooled to -196°C. Journal of Plant Physiology, 137: 465-470.
- Seitz U., Reinhard E. (1987). Growth and ginsenoside patterns of cryopreserved *Panax ginseng* cell cultures. Journal of Plant Physiology, 131: 215-223.
- Warmund M. R., Starbuck C. J., George M. F. (1989). Freezing tolerance of tissue cultured *Rubus* plants. Fruit Varieties Journal, 43: 109-114.
- Yamada T., Sakai A., Matsumura T., Higuchi S. (1991). Cryopreservation of apical meristems of white clover (*Trifolium repens* L.) by vitrification. Plant Science, 78: 81-87.