

**IDENTIFICATION AND QUANTITATIVE ANALYSIS OF GERMINATION INHIBITORS
IN THE PERICARP OF *CYCLOCARYA PALIURUS* (BATAL) ILJINSKAJA**

Xulan Shang*, Xizeng Xu, and Shengzuo Fang

College of Forest Resources and Environment, Nanjing Forestry University, Nanjing 210037, China,

*Fax: + 86 25 85428664, *E-mail: sxl66@sina.com

REFERENCES

- BENECH-ARNOLD R., GUALANO N., LEYMARIE J., COME D., CORBINEAU F. (2006). Hypoxia interferes with ABA metabolism and increases ABA sensitivity in embryos of dormant barley grains. *Journal of Experimental Botany*, 57: 1423-1430.
- BRIGGS C. L., MORRIS E. C., ASHFORD A. E. (2005). Investigations into seed dormancy in *Grevillea linearifolia*, *G. buxifolia* and *G. sericea*: Anatomy and histochemistry of the seed coat. *Annals of Botany*, 96: 965-980.
- FANG S. Z., WANG J. Y., WEI Z. Y., ZHU Z. X. (2006). Methods to break seed dormancy in *Cyclocarya paliurus* (Batal) Iljinskaja. *Scientia Horticulturae*, 110: 305-309.
- FANG S. Z., YANG W. X., CHU, X. L., SHANG X. L., SHE C. Q., FU X. X. (2011). Provenance and temporal variations in selected flavonoids in leaves of *Cyclocarya paliurus*. *Food Chemistry*, 124: 1382-1386.
- GALLAGHER R. S., ANANTH R., GRANGER K., BRADLEY B., ANDERSON J. V., FUERST E. P. (2010). Phenolic and short-chained aliphatic organic acid constituents of wild oat (*Avena fatua* L.) seeds. *Journal of Agricultural and Food Chemistry*, 58: 218-225.
- GATFORD K. T., EASTWOOD R. F., HALLORAN G. M. (2002). Germination inhibitors in bracts surrounding the grain of *Triticum tauschii*. *Functional Plant Biology*, 29: 881-890.
- GRANGER K. L., GALLAGHER R. S., FUERST E. P., ALLDREDGE J. R. (2011). Comparison of seed phenolic extraction and assay methods. *Methods in Ecology and Evolution*, 2: 691-698.
- HURA T., DUBERT F., DABKOWSKA T., STUPNICKA-RODZYNKIEWICZ E., STOKLOSA A., LEPIARCZYK A. (2006). Quantitative analysis of phenolics in selected crop species and biological activity of these compounds evaluated by sensitivity of *Echinochloa crus-galli*. *Acta Physiologiae Plantarum*, 28: 537-545.
- KALINOVA J., VRCHOTOVA N., TRISKA J. (2007). Exudation of allelopathic substances in buckwheat (*Fagopyrum esculentum* Moench). *Journal of Agricultural and Food Chemistry*, 55: 6453-6459.
- KONG L. G., WANG F. H., SI J. S., FENG B., LI S. D. (2008). Water-soluble phenolic compounds in the coat control germination and peroxidase reactivation in *Triticum aestivum* seeds. *Plant Growth Regulation*, 56: 275-283.
- KURIHARA H., ASAMI S., SHIBATA H., FUKAMI H., TANAKA T. (2003). Hypolipemic effect of *Cyclocarya paliurus* (Batal) Iljinskaja in lipid-loaded mice. *Biological and Pharmaceutical Bulletin*, 26: 383-385.
- KUSHIMA M., KAKUTA H., KOSEMURA S., YAMAMURA S., YAMADA K., YOKOTANI-TOMITA K., HASEGAWA K. (1998). An allelopathic substance exuded from germinating watermelon seeds. *Plant Growth Regulation*, 25: 1-4.
- MUSCOLO A., PANUCCIO M. R., SIDARI M. (2001). The effect of phenols on respiratory enzymes in seed germination - respiratory enzyme activities during germination of *Pinus laricio* seeds treated with phenols extracted from different forest soils. *Plant Growth Regulation*, 35: 31-35.
- SAMPIETRO D. A., VATTUONE M. A., ISLA M. A. (2006). Plant growth inhibitors isolated from sugarcane (*Saccharum officinarum*) Straw. *Journal of Plant Physiology*, 163: 837- 846.
- SHANG X. L., XU X. Z., FANG S. Z. (2011). Seed dormancy mechanism of *Cyclocarya paliurus*. *Scientia Silvae Sinicae*, 47: 68-74 (in Chinese).
- SHE C. Q., FANG S. Z., YANG W. X. (2008). Geographical variation of morphologic characteristics of *Cyclocarya paliurus* seeds. *Journal of Nanjing Forestry University (Natural Sciences Edition)*, 32: 63-66 (in Chinese).
- SOLTANI A. (2004). Phenolics in the seed coat of oriental beech (*Fagus orientalis* Lipsky) as an autotoxic factor. *Proceedings of the Fourth International Iran and Russia Conference*: 1048-1055.
- WEIDNER S., AMAROWICZ R., KARAMAC M., FRACZEK E. (2000). Changes in endogenous phenolic acids during development of *Secale cereale* caryopses and after dehydration treatment of unripe rye grains. *Plant Physiology and Biochemistry*, 38: 595-602.
- WEIDNER S., KRUPA U., AMAROWICZ R., KARAMAC M., ABE S. (2002). Phenolic compounds in embryos of triticale caryopses at different stages of development and maturation in normal environment and after dehydration treatment. *Euphytica*, 126: 115-122.
- WROBEL M., KARAMAC M., AMAROWICZ R., FRQCZEK E., WEIDNER S. (2005). Metabolism of phenolic compounds in *Vitis riparia* seeds during stratification and during germination under optimal and low temperature stress conditions. *Acta Physiologiae Plantarum*, 27: 313-320.
- XIAO W., LI N., LI X. (2008). Isolation and identification of organic acids from pericarp of *Capparis spinosa* L. *Journal of Shenyang Pharmaceutical University*, 25: 790-792 (in Chinese).
- XIE J. H., XIE M. Y., NIE S. P., SHEN M. Y., WANG Y. X., LI C. (2010). Isolation, chemical composition and antioxidant activities of a water-soluble polysaccharide from *Cyclocarya paliurus* (Batal.) Iljinskaja. *Food Chemistry*, 119: 1626-1632.
- YANG W. X., FANG S. Z. (2005). The bioassay of the methanol extract from *Cyclocarya paliurus* seed coat. *Journal of Plant Resources and Environment*, 14: 11-14 (in Chinese).
- ZHANG X. Q., YE W. C., YIN Z. Q., ZHANG Z. H., ZHAO S. X. (2005). Study of chemical constituents of *Cyclocarya paliurus*. *China Journal of Chinese Materia Medica*, 30: 91-92 (in Chinese).