

**PLANT REGENERATION FROM LIQUID CALLUS CULTURES DERIVED
FROM MATURE SEEDS OF PAMPAS GRASS
[*CORTADERIA SELLOANA* (SCHULT. & SCHULT. F.) ASCH. & GRAEBN.]**

**Kazuki Shimomae¹, Hiroyuki Niimi¹, Dong Poh Chin¹, Raham Sher Khan^{1,2}, Yuanxue Lu³,
Yoichi Ogawa⁴, and Masahiro Mii^{1,4*}**

¹Laboratory of Plant Cell Technology, Graduate School of Horticulture, Chiba University,
648 Matsudo, Matsudo, Chiba 271-8510, Japan

*Fax +81-47-308-8852, E-mail: miim@faculty.chiba-u.jp

²Department of Biotechnology, Abdul Wali Khan University Mardan, Pakistan

³Key Laboratory of Economic Plants and Biotechnology, Kunming Institute of Botany,
Chinese Academy of Sciences, 132 Lanhei Road, Kunming, Yunnan, 650204, China

⁴Center for Environment, Health and Field Sciences, Chiba University,
6-2-1 Kashiwanoha, Kashiwa, Chiba 277-0882, Japan

REFERENCES

- AGUADO-SANTACRUZ G. A., CABRERA-PONCE J. L., OLALDE-PORTUGAL V., SANCHEZ-GONZALEZ M. R., MÁRQUEZ-GUZMÁN J., HERRERA-ESTRELLA L. (2001). Tissue culture and plant regeneration of blue grama grass, *Bouteloua gracilis* (H.B.K.) Lag. ex Steud. In *In Vitro Cellular & Developmental Biology-Plant*, 37: 182-189.
- AKUTSU M., SATO H. (2002). Induction of proembryos in liquid culture increases the efficiency of plant regeneration from *Alstroemeria* calli. *Plant Science*, 163: 475-479.
- CONNOR H. E., EDGAR E. (1974). Names and types of *Cortaderia* Stapf (Graminae). *Taxon*, 23: 595-605.
- HORNBACK B. (1994). In praise of pampas grass. *Pacific Horticulture*, 56: 47-52.
- JOYCE P., KUWAHATA M., TURNER N., LAKSHMANAN P. (2010). Selection system and co-cultivation medium are important determinants of *Agrobacterium*-mediated transformation of sugarcane. *Plant Cell Reports*, 29: 173-183.
- MADISON J. (1992). Pampas grasses: one a weed and one a garden queen. *Pacific Horticulture*, 53: 48-53.
- MAURY S., TRAP-GENTIL M. V., HÉBRARD C., WEYENS G., DELAUNAY A., BARNES S., LEFEBVRE M., JOSEPH C. (2012). Genic DNA methylation changes during in vitro organogenesis: Organ specificity and conservation between parental lines of epialleles. *Physiologia Plantarum*, 146: 321-335.
- MURASHIGE T., SKOOG F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- OCHATT S. J., MOUSSET-D'ÉCLAS C., RANCILLAC M. (2000). Fertile pea plants regenerate from protoplasts when calluses have not undergone endoreduplication. *Plant Science*, 156: 177-183.
- OZAWA K., TAKAIWA F. (2010). Highly efficient *Agrobacterium*-mediated transformation of suspension-cultured cell cluster of rice (*Oryza sativa* L.). *Plant Science*, 179: 333-337.
- ROBACKER C. (1995). Long-term shoot regeneration from pampas grass (*Cortaderia selloana* Schult.) through manipulation of growth regulators *in vitro*. *Plant Cell Reports*, 14: 689-693.
- ROBACKER C. (1995). Long-term shoot regeneration from pampas grass (*Cortaderia selloana* Schult.) through manipulation of growth regulators *in vitro*. *Plant Cell Reports*, 14: 689-693.
- ROBACKER C. D., CORLEY W. L. (1992). Plant regeneration of pampas grass from immature inflorescences cultured *in vitro*. *Hort-Science*, 27: 841-843.
- ŞENER O., CAN E., ARSLAN M., ÇELIKTAS N. (2008). Effects of genotype and picloram concentrations on callus induction and plant regeneration from immature inflorescence of spring barley cultivars (*Hordeum vulgare* L.). *Biotechnology & Biotechnological Equipment*, 22: 915-920.
- SEO M., TAKAHARA M., TAKAMIZO T. (2010). Optimization of culture conditions for plant regeneration of *Panicum* spp. through somatic embryogenesis. *Grassland Science*, 56: 6-12.
- SHIBA T., MII M. (2005). Visual selection and maintenance of cell lines with high plant regeneration ability and low level in *Dianthus acicularis* by monitoring with flow cytometry analysis. *Plant Cell Reports*, 24: 572-580.
- SHIMIZU K., HASHIMOTO M., HASHIMOTO F., SAKATA Y. (2003). Plant regeneration from suspension cultures in Japanese morning glory (*Ipomoea nil* (L.) Roth.). *Journal of the Japanese Society for Horticulture Science*, 72: 409-414.
- SINGH R. J. (1986). Chromosomal variation in immature embryo derived calluses of barley (*Hordeum vulgare* L.). *Theoretical and Applied Genetics*, 72: 710-716.
- YUAN X., WANG Z., LIU J., SHE J. (2009). Development of a plant regeneration system from seed-derived calluses of centipedegrass [*Eremochloa ophiuroides* (Munro.) Hack.]. *Scientia Horticulturae*, 120: 96-100.
- ZHANG Q. X., SUN Y, HU H. K., CHEN B., HONG C. T., GUO H. P., PAN Y. H., ZHENG B. S. (2012). Micropropagation and plant regeneration from embryogenic callus of *Miscanthus sinensis*. *In Vitro Cellular & Developmental Biology-Plant*, 48: 50-57.

