

## DOES SHOOT ANATOMICAL HETEROGENEITY INFLUENCE *EX VITRO* ADVENTITIOUS ROOT FORMATION IN RHODODENDRON MICROCUTTINGS?

Kristine Dokane<sup>1\*</sup>, Dace Megre<sup>1,3</sup>, Madara Lazdane<sup>2</sup>, and Uldis Kondratovics<sup>1</sup>

<sup>1</sup>Faculty of Biology, University of Latvia, 4 Kronvalda Boulv., 1586 Riga, Latvia,

\*Fax: + 371-67034862, \*E-mail: kristine.dokane@lu.lv

<sup>2</sup>Plant Biology Laboratory, Botanical Garden, University of Latvia, 2 Kandavas str., 1083 Riga, Latvia

<sup>3</sup>National Botanical Garden, 1 Miera str., 2169 Salaspils, Latvia

### REFERENCES

- ALTAMURA M. M. (1996). Root histogenesis in herbaceous and woody explants cultured *in vitro*. A critical review. *Agronomie*, 16: 589-602.
- ANDERSON W. C. (1984). A revised tissue culture medium for shoot multiplication of rhododendron. *Journal of the American Society for Horticultural Sciences*, 109: 343-347.
- BALLESTER A., SAN-JOSE M. C., VIDAL N., FERNANDEZ-LORENZO J. L., VIEITEZ A. M. (1999). Anatomical and biochemical events during *in vitro* rooting of microcuttings from juvenile and mature phases of chestnut. *Annals of Botany*, 83: 619-629.
- BRAUNE W., LEMAN A., TAUBERT H. (1999). *Pflanzenanatomisches praktikum I*. Spektrum Akademischer Verlag Heidelberg, Berlin, 368 pp.
- DE KLERK G.-J. (2002). Rooting of microcuttings: theory and practice. *In Vitro Cellular & Developmental Biology-Plant*, 38: 415-422.
- DE KLERK G.-J., VAN DER KRIEKEN W., DE JONG J. C. (1999). The formation of Adventitious roots: new concepts, new possibilities. *In Vitro Cellular & Developmental Biology-Plant*, 35: 189-199.
- DONNELLY D. J., VIDAVER W. E., LEE K. Y. (1985). The anatomy of tissue cultured red raspberry prior to and after transfer to soil. *Plant Cell, Tissue Organ Culture*, 4: 43-50.
- ERMEL F. F., VIZOSO S., CHARPENTIER J.-P., JAY-ALLEMAND CH., CATESSON A.-M., COUÉE I. (2000). Mechanisms of primordium formation during root development from walnut cotyledon explants. *Planta*, 211: 563-574.
- FAHN A. (1990). *Plant Anatomy*. Pergamon Press, Oxford, 588 pp.
- GASPAR T., KEVERS C., HAUSMAN J. F., RIPETTI V. (1994). Peroxidase activity and endogenous free auxin during adventitious root formation. *In: Lumsden P. J., Nicholas J. R., Davies W. J. (Eds). Physiology, Growth and Development of Plants in Culture*, Kluwer Academic Publishers: 289-298.
- GEORGE E. F., HALL M. A., DE KLERK G.-J. (2008). *Plant Propagation by Tissue Culture 3<sup>rd</sup> Edition*. Springer, The Netherlands, 501 pp.
- HARBAGE J. F., STIMART D. P., EVERT R. F. (1993). Anatomy of adventitious root formation in microcuttings of *Malus domestica* Borkh. 'Gala'. *Journal of the American Society for Horticultural Science*, 118: 680-688.
- HARTMANN H. T., KESTER D. E., DAVIES F. T., GENEVE R. L. (2011). *Plant Propagation Principles and Practices 8<sup>th</sup> edition*. Prentice Hall, New Jersey, 880 pp.
- HATZILAZAROU S. P., SYROS T. D., YUPSANIS T. A., BOSABALIDIS A. M., ECONOMOU A. S. (2006). Peroxidases, lignin and anatomy during *in vitro* and *ex vitro* rooting of gardenia (*Gardenia jasminoides* Ellis) microshoots. *Journal of Plant Physiology*, 163: 827-836.
- ILIEV I., KITIN P., FUNADA R. (2001). Morphological and anatomical study on *in vitro* root formation of silver birch (*Betula pendula* Roth.). *Propagation of Ornamental Plants*, 1: 10-19.
- KONDRATOVIČS U., MEGRE D. (1999). Anatomische eigenarten der adventivwurzelbildung von Rhododendronstecklingen während der bewurzelung. *Rhododendron und immergrüne Laubgehölze, Jahrbuch*: 72-85.
- LEROUX O. (2012). Collenchyma: a versatile mechanical tissue with dynamic cell walls. *Annals of Botany*, 110: 1083-1098.
- MEGRE D., DOKANE K., KONDRATOVIČS U. (2011). Can changes in starch content and peroxidase activity be used as rooting phase markers for rhododendron leaf bud cuttings? *Acta Biologica Cracoviensia, Series Botanica*, 53: 74-79.
- MILLÁN-OROZCO L., CORREDOIRA E., SAN-JOSÉ M. C. (2011). *In vitro* rhizogenesis: histoanatomy of *Cedrela odorata* (Meliaceae) microcuttings. *International Journal of Tropical Biology*, 59: 447-453.
- REVERBERI M., FALASCA G., LAURI P., CABONI E., ALTAMURA M. M. (2001). Indoleacetic acid induces xylem formation instead of rooting in walnut (*Juglans regia* L.) microcuttings. *Plant Biosystems*, 135: 71-77.
- RUZIN S. E. (1999). *Plant Microtechnique and Microscopy*. Oxford University Press, New York, 322 pp.
- SAN-JOSÉ M. C., VIDAL A., BALLESTER A. (1992). Anatomical and biochemical changes during root formation in oak and apple shoots cultured *in vitro*. *Agronomie*, 12: 767-774.
- SAN-JOSÉ M. C., ROMERO L., JANEIRO L. V. (2012). Effect of indole-3-butyric acid on root formation in *Alnus glutinosa* microcuttings. *Silva Fennica*, 46: 643-654.
- YEREMIN V. M., BOIKO V. I. (1998). Bark anatomy of the stem in some species of the Ericaceae. *Botanicheskii Zhurnal*, 83 (8): 1-14 (in Russian).
- ZHOU J., WU H., COLLET G. F. (1992). Histological study of initiation and development *in vitro* of adventitious roots in minicuttings of apple rootstocks of M26 and EMLA9. *Physiologia Plantarum*, 84: 433-440.