Propagation of Ornamental Plants
Vol. 16, № 3, 2016: 79-83

EFFECT OF *META*-TOPOLIN ON ADVENTITIOUS SHOOT REGENERATION IN *BEGONIA SEMIPARIETALIS* Y. LIU, S. M. KU & C. I PENG

Tsai-Yu Chung[1], Li-Peng Hsu[1], and Wei-Hsin Hu[2]*

[1]Department of Bioresources, Dayeh University, 168 University Rd. Dacun, 515 Changhua, Taiwan,
[2]Department of Biology, National Museum of Natural Science, 1 Guancian Rd., 404 Taichung, Taiwan
*Fax: + 886-4-23285326, *E-mail: bohu@mail.nmns.edu.tw

REFERENCES

Amoo S. O., Finnie J. F., Van Staden J. (2011). The role of *meta*-topolins in alleviating micropropagation problems. Plant Growth Regulation, 63: 197-206.

Aremu A. O., Bairu M. W., Dolezal K., Finnie J. F., Van Staden J. (2012). Topolins: a panacea to plant tissue culture challenges? Plant Cell, Tissue and Organ Culture, 108: 1-16.

Awal A., Ahmed A. B. A., Taha R. M., Yaacob J. S., Mohajer S. (2013). Effect of adenine, sucrose and plant growth regulators on the indirect organogenesis and on *in vitro* flowering in *Begonia* × *hiemalis* Fotsch. Australian Journal of Crop Science, 7: 691-698.

Bairu M. W., Stirk W. A., Dolezal K., Van Staden J. (2007). Optimizing the micropropagation protocol for the endangered *Aloe polyphylla*: can *meta*-topolin and its derivatives serve as replacement for benzyladenine and zeatin? Plant Cell, Tissue and Organ Culture, 90: 15-23.

Burritt D. J., Leung D. W. M. (2003). Adventitious shoot regeneration from *Begonia* × *erythrophylla* petiole sections is developmentally sensitive to light quality. Physiologia Plantarum, 118: 289-296.

Chung K. F., Leong W. C., Rubite R. R., Repin R., Kiew R., Liu Y., Peng C. I (2014). Phylogenetic analyses of *Begonia* sect. *Coelocentrum* and allied limestone species of China shed light on the evolution of Sino-Vietnamese karst flora. Botanical Studies, 55: 15 pp. DOI: 10.1186/1999-3110-55-1.

Godo T., Lu Y. X., Li J. X., Guan K. Y. (2008). Comparisons of response for plant growth regulators in tissue culture of *Begonia* native to Yunnan, China. Bulletin of the Botanical Garden of Toyama, 13: 41-46.

Hu W. H., Peng C. I, Liaw S. I. (2012). Adventitious bud regeneration from lamina culture of *Begonia parvula* H. Lév., and Vaniot. Seed and Nursery, 14: 21-32 (in Chinese).

Kaviani B., Hashemabadi D., Khodabakhsh H., Onsinejad R., Ansari M. H., Haghighat N. (2015). Micropropagation of *Begonia rex* Putz. by 6-benzyladenine and α-naphthalene acetic acid. International Journal of Biosciences, 6: 8-15.

Köszeghi S., Bereczki C., Balog A., Benedek K. (2014). Comparing the effects of benzyladenine and *meta*-Topolin on Sweet Basil (*Ocimum basilicum*) micropropagation. Notulae Scientia Biologicae, 6: 422-427.

Ku S. M., Liu Y., Peng C. I (2006). Four new species of *Begonia* sect. *Coelocentrum* (Begoniaceae) from limestone areas in Guangxi, China. Botanical Studies, 47: 207-222.

Kubalakova M., Strnad M. (1992). The effects of aromatic cytokinins on micropropogation and regeneration of sugar beet *in vitro*. Biologia Plantarum, 34: 578-579.

Kumaria S., Kehie M., Bhowmik S. S. D., Singh M., Tandon P. (2012). *In vitro* regeneration of *Begonia rubrovenia* var. *meisneri* C.B. Clarke – a rare and endemic ornamental plant of Meghalaya, India. Indian Journal of Biotechnology, 11: 300-303.

Mendi Y. Y., Curuk P., Kocaman E., Unek C., Eldogan S., Gencel G., Cetiner S. (2009). Regeneration of begonia plantlets by direct organogenesis. African Journal of Biotechnology, 8: 1860-1863.

Murashige T., Skoog F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiologia Plantarum, 15: 473-497.

Nada S., Chennareddy S., Goldman S. (2011). Direct shoot bud differentiation and plantlet regeneration from leaf and petiole explants of *Begonia tuberhybrida.* HortScience, 46: 759-764.

Nakano M., Niimi Y., Kobayashi D., Watanabe A. (1999). Adventitious shoot regeneration and micropropagation of hybrid tuberous begonia (*Begonia* × *tuberhybrida* Voss). Scientia Horticulturae, 79: 245-251.

Nakano M., Nagai M., Tanaka S., Nakata M., Godo T. (2005). Adventitious shoot regeneration and micropropagation of the Japanese endangered *Hylotelephium sieboldii* (Sweet ex Hook.) H. Ohba and *H. sieboldii* var. *ettyuense* (Tomida) H. Ohba. Plant Biotechnology, 22: 221-224.

Nhut D. T., Hai N. T., Phan M. X. (2010). A highly efficient protocol for micropropagation of *Begonia tuberous. In*: Jain S. M., Ochatt S. J. (Eds). Protocols for *in vitro* propagation of ornamental plants. Humana Press, USA: 15-20.

Peng C. I, Ku S. M., Kono Y., Chung K. F., Liu Y. (2008). Two new species of *Begonia* (sect. *Coelocentrum*, Begoniaceae) from limestone areas in Guangxi, China: *B. arachnoidea* and *B. subcoriacea.* Botanical Studies, 49: 405-418.

Peng C. I, Ku S. M., Yang H. A., Leong W .C., Liu Y., Nguyen T. H., Kono Y., Chung K. F. (2014). Two new species of *Begonia* sect. *Coelocentrum*, *B. guixiensis* and *B. longa*, from Sino-Vietnamese limestone karsts. Botanical Studies, 55: 12 pp. DOI: 1186/s40529-014-0052-8.

Peng C. I, Lin C. W., Yang H. A., Kono Y., Nguyen T. H. (2015). Six new species of *Begonia* (Begoniaceae) from limestone areas in Northern Vietnam. Botanical Studies, 56: 23 pp. DOI: 10.1186/s40529-015-0089-3.

Van Staden J., Zazimalova E., George E. F. (2008). Chapter 6: Plant growth regulators, II: Introduction; cytokinins, their analogues and inhibitors. *In*: George E. F., Hall M. A., de Klerk G. J. (Eds). Plant Propagation by tissue culture, 3th ed., vol. 1, Springer, Dordrecht, The Netherlands: 205-226.

Werbrouck S. P. O., Van der Jeugt B., Dewitte W., Prinsen E., Van Onckelen H. A., Debergh P. C. (1995). The metabolism of benzyladenine in *Spathiphyllum floribundum* 'Schott Petite' in relation to acclimatization problems. Plant Cell Reports, 14: 662-665.

Werbrouck S. P. O., Strnad M., Van Onckelen H. A., Debergh P. C. (1996). *Meta*-topolin, an alternative to benzyladenine in tissue culture? Physiologia Plantarum, 81: 291-297.