

**IN VITRO REGENERATION AND PROPAGATION OF  
HYDRANGEA MACROPHYLLA THUNB. 'NACHTIGALL'**

**Antje Doil<sup>1</sup>\*, Rui Zhang<sup>2</sup>, Annegret Schum<sup>3</sup>, Margrethe Serek<sup>2</sup>, and Traud Winkelmann<sup>1</sup>**

<sup>1</sup>University of Applied Sciences Weihenstephan and Research Station for Horticulture, Am Staudengarten 8, D-85354 Freising, Germany, \*Fax: + 49 8161 715106, \*E-mail: antje.doil@fh-weihenstephan.de

<sup>2</sup>Gottfried Wilhelm Leibniz University of Hannover, Institute of Ornamental and Woody Plant Science, Floriculture Section, 2 Herrenhaeuser str., D-30419 Hannover, Germany

<sup>3</sup>Federal Research Centre for Cultivated Plants - Julius Kuehn Institute, Institute of Resistance Research and Stress Tolerance, Rudolf-Schick-Platz 3, D-18190 Gross Luesewitz, Germany

**REFERENCES**

- Boccon-Gibod J., Billard C., Maltete S. (2000). *In vitro* regeneration system of *Hydrangea macrophylla* plantlets from leaves and internodes. *Acta Horticulturae*, 508: 229-231.
- Gamborg O. L., Miller R. A., Ojima K. (1968). Nutrient requirement suspension cultures of soybean root cells. *Experimental Cell Research*, 50: 151-158.
- Lorenzo J. C., Gonzales B. L., Escalona M., Teisson C., Espinosa P., Borroto C. (1998). Sugarcane shoot formation in an improved temporary immersion system. *Plant Cell, Tissue and Organ Culture*, 54: 197-200.
- Murashige T., Skoog F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 18: 100-127.
- Preece J. E., Ledbetter D. I. (2003). The influence of thidiazuron on *in vitro* shoot proliferation of oakleaf hydrangea (*Hydrangea quercifolia* Bartr.). *Acta Horticulturae*, 625: 233-236.