

**ADVENTITIOUS SHOOT FORMATION FROM LEAF EXPLANTS
OF OLEASTER (*ELAEAGNUS ANGUSTIFOLIA* L.)**

Omid Karami*, Ali Deljou, and Khosro Piri

Department of Biotechnology, Faculty of Agriculture, Bu-Ali Sina University, Hamadan, Iran

*Fax: + 8114227021, *E-mail: hiva@basu.ac.ir

REFERENCES

- Blakesey D. (1991). Uptake and metabolism of 6-benzyladenine in shoot proliferation of *Musa* and *Rhododendron*. *Plant Cell, Tissue and Organ Culture*, 25: 69-74.
- Burne C. (2006). Tissue culture for gene transfer. *Farming Ahead*, 172: 45.
- Economou A. S., Maloupa E. M. (1995). Regeneration of *Elaeagnus angustifolia* from leaf segments of *in vitro*-derived shoots. *Plant Cell, Tissue and Organ Culture*, 40: 285-288.
- Economou A. S., Spanoudaki M. J. (1988). Regeneration *in vitro* of oleaster (*Elaeagnus angustifolia* L.) from shoot tips of mature tree. *Acta Horticulturae*, 227: 363-368.
- Hicks G. S. (1994). Shoot induction and organogenesis *in vitro*: a developmental perspective. *In Vitro Cellular and Developmental Biology-Plant*, 30: 10-15.
- Iriondo M., Delaiglesia M., Perez (1995). Micropropagation of *Elaeagnus angustifolia* from mature trees. *Tree Physiology*, 15: 691-693.
- Kamínek M. (1992). Progress in cytokinin research. *Trends in Biotechnology*, 10: 159-162.
- Lin C.-S., Chih Lin C., Chang W.-C. (2004). Effect of thidiazuron on vegetative tissue-derived somatic embryogenesis and flowering of bamboo *Bambusa edulis*. *Plant Cell, Tissue and Organ Culture*, 76: 75-82
- Lu C. Y. (1993). The use of thidiazuron in tissue culture. *In Vitro Cellular and Developmental Biology-Plant*, 29: 92-96
- Martin K. P. (2004). Benzyladenine induced somatic embryogenesis and plant regeneration of *Leptadenia reticulata*. *Biologia Plantarum*, 48: 285-288.
- Murashige T., Skoog, F. A. (1962). Revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 154: 73-479.
- Murthy B. N. S., Murch S. J., Saxena P. K. (1995). Thidiazuron-induced somatic embryogenesis in intact seedlings of peanut (*Arachis hypogaea* L.): endogenous growth regulator levels and significance of cotyledons. *Physiologia Plantarum*, 94: 268-276.
- Schimuling T. (2002). New insights into the functions of cytokinins in plant development. *Plant Growth Regulation*, 21 : 40-49.
- Srinivasan M., Nnchiappan V., Rajaskharna R. (2006). Potential application of urea-derived herbicides as cytokinins in plant tissue culture. *Journal of Bioscience*, 31: 599-605.
- Vetvicka V., Matousova V. (1991). *Arboles y arbustos*. Susaeta, Madrid, Spain, 311 p.
- Victor J. M. R., Murch S. J., Krishnaraj S., Saxena P. K. (1999). Somatic embryogenesis and organogenesis in peanut: The role of thidiazuron and N⁶-benzylaminopurine in the induction of plant morphogenesis. *Plant Growth Regulation*, 28: 9-15.
- Visser C., Fletcher R. A., Saxena P. K. (1995). Thidiazuron stimulates expansion and greening in cucumber cotyledons. *Physiology & Molecular Biology of Plants*. 1: 21-25.